

PhenoMaster



Publications for

PhenoMaster

2015

- **Finan B, Yang B, Ottaway N, Smiley DL, Ma T, Clemmensen C, Chabenne J, Zhang L, Habegger KM, Fischer K, Campbell JE, Sandoval D, Seeley RJ, Bleicher K, Uhles S, Riboulet W, Funk J, Hertel C, Belli S, Sebokova E, Conde-Knape K, Konkar A, Drucker DJ, Gelfanov V, Pfluger PT, Müller TD, Perez-Tilve D, DiMarchi RD, Tschöp MH.** A rationally designed monomeric peptide triagonist corrects obesity and diabetes in rodents. *Nat Med* 2015; 21(1):27-36
<http://www.ncbi.nlm.nih.gov/pubmed/25485909>
- **Sudakov SK, Bogdanova NG, Kolpakov AA.** Method for determining the level of food motivation in rats. *Exp Biol Med* 2015; 158(3):401-3
<http://www.ncbi.nlm.nih.gov/pubmed/25573375>
- **Sudakov SK, Nazarova GA, Alekseeva EV.** Changes in feeding behavior, locomotor activity, and metabolism in rats upon modulation of opioid receptors in the gastrointestinal tract. *Bull Exp Biol Med* 2015; 156(4):423-5
<http://www.ncbi.nlm.nih.gov/pubmed/24771418>
- **Tang C, Ahmed K, Gille A, Lu S, Gröne HJ, Tunaru S, Offermanns S.** Loss of FFA2 and FFA3 increases insulin secretion and improves glucose tolerance in type 2 diabetes. *Nat Med* 2015; Epub ahead of print
<http://www.ncbi.nlm.nih.gov/pubmed/25581519>
- **Tranberg B, Madsen AN, Hansen AK, Helligren LI.** Whey-reduced weight gain is associated with a temporary growth reduction in young mice fed a high-fat diet. *J Nutr Biochem* 2015; 26(1):9-15
<http://www.ncbi.nlm.nih.gov/pubmed/25315863>

2014

- **Adamovich Y, Rousso-Noori L, Zwihaft Z, Neufeld-Cohen A, Golik M, Kraut-Cohen J, Wang M, Han X, Asher G.** Circadian clocks and feeding time regulate the oscillations and levels of hepatic triglycerides. *Cell Metab* 2014; 19(2):319-30
<http://www.ncbi.nlm.nih.gov/pubmed/24506873>
- **Anderson RI, Becker HC, Adams BL, Jesudason CD, Rorick-Kehn LM.** Orexin-1 and orexin-2 receptor antagonists reduce ethanol self-administration in high-drinking rodent models. *Front Neurosci* 2014; 8:33
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3933945/>
- **Bashkatova V, Sudakov S, Nazarova G, Alexeeva E.** The Action of Chronic Nicotine on the Effects of Ethanol on Anxiety, Locomotion and Metabolism and the Feeding and Drinking Behaviors of Rats. *Pharmacology & Pharmacy* 2014; 5: 1077-84
http://www.scirp.org/journal/PaperInformation.aspx?paperID=50993#.VHLuhPmG_w8

- **Bake T, Murphy M, Morgan DG, Mercer JG.** Large, binge-type meals of high fat diet change feeding behaviour and entrain food anticipatory activity in mice. *Appetite* 2014; 77:62-73
<http://www.ncbi.nlm.nih.gov/pubmed/24631639>
- **Cansell C, Castel J, Denis RG, Rouch C, Delbes AS, Martinez S, Mestivier D, Finan B, Maldonado-Aviles JG, Rijnsburger M, Tschöp MH, Dileone RJ, Eckel RH, la Fleur SE, Magnan C, Hnasko TS, Luquet S.** Dietary triglycerides act on mesolimbic structures to regulate the rewarding and motivational aspects of feeding. *Mol Psychiatry* 2014; Epub ahead of print
<http://www.ncbi.nlm.nih.gov/pubmed/24732670>
- **Castelhano-Carlos M, Costa PS, Russig H, Sousa N.** PhenoWorld: a new paradigm to screen rodent behavior. *Transl Psychiatry* 2014; 4:e399
<http://www.nature.com/tp/journal/v4/n6/full/tp201440a.html>
- **Clemens LE, Jansson EK, Portal E, Riess O, Nguyen HP.** A behavioral comparison of the common laboratory rat strains Lister Hooded, Lewis, Fischer 344 and Wistar in an automated homecage system. *Genes Brain Behav* 2014; 13(3):305-21
<http://www.ncbi.nlm.nih.gov/pubmed/24119005>
- **Collden G, Balland E, Parkash J, Caron E, Langlet F, Prevot V, Bouret SG.** Neonatal overnutrition causes early alterations in the central response to peripheral ghrelin. *Mol Metab* 2014; Epub ahead of print
<http://www.sciencedirect.com/science/article/pii/S2212877814001781>
- **Constantinou C, Mpatsoulis D, Natsos A, Petropoulou PI, Zvintzou E, Traish AM, Voshol PJ, Karagiannides I, Kypreos KE.** The low density lipoprotein receptor modulates the effects of hypogonadism on diet-induced obesity and related metabolic perturbations. *J Lipid Res* 2014; 55(7):1434-47
<http://www.ncbi.nlm.nih.gov/pubmed/24837748>
- **Diep TA, Madsen AN, Krogh-Hansen S, Al-Shahwani M, Al-Sabagh L, Holst B, Hansen HS.** Dietary Non-Esterified Oleic Acid Decreases the Jejunal Levels of Anorectic N-Acylethanolamines. *PLoS One* 2014; 9(6):e100365
<http://www.ncbi.nlm.nih.gov/pubmed/24959837>
- **Duivenvoorde LP, van Schothorst EM, Derosus D, van der Stelt I, Masania J, Rabbani N, Thornalley PJ, Keijer J.** Oxygen restriction as challenge test reveals early high-fat-diet-induced changes in glucose and lipid metabolism. *Pflugers Arch* 2014; Epub ahead of print
<http://www.ncbi.nlm.nih.gov/pubmed/24974902>
- **Farzi A, Reichmann F, Meinitzer A, Mayerhofer R, Jain P, Hassan AM, Fröhlich EE, Wagner K, Painsipp E, Rinner B, Holzer P.** Synergistic effects of NOD1 or NOD2 and TLR4 activation on mouse sickness behavior in relation to immune and brain activity markers. *Brain Behav Immun* 2014; Epub ahead of print
<http://www.ncbi.nlm.nih.gov/pubmed/25218901>
- **Ikels K, Kuschel S, Fischer J, Kaisers W, Eberhard D, Rütther U.** FTO Is a Relevant Factor for the Development of the Metabolic Syndrome in Mice. *PLoS One* 2014 9(8):e105349
<http://www.ncbi.nlm.nih.gov/pubmed/25144618>

- **Jodayree S, Patterson ZR, MacKay H, Abizaid AB, Tsopmo A.** Blood and Liver Antioxidant Capacity of Mice Fed High Fat Diet Supplemented with Digested Oat Bran Proteins. *International Journal of Food Science and Nutrition Engineering* 2014; 4(1):9-14
<http://article.sapub.org/10.5923.j.food.20140401.02.html>
- **Kasza I, Suh Y, Wollny D, Clark RJ, Roopra A, Colman RJ, MacDougald OA, Shedd TA, Nelson DW, Yen MI, Yen CL, Alexander CM.** Syndecan-1 Is Required to Maintain Intradermal Fat and Prevent Cold Stress. *PLoS Genet* 2014; 10(8):e1004514
<http://www.ncbi.nlm.nih.gov/pubmed/25101993>
- **Keijer J, Hoevenaars FP, Nieuwenhuizen A, van Schothorst EM.** Nutrigenomics of body weight regulation: a rationale for careful dissection of individual contributors. *Nutrients* 2014; 6(10): 4531-51
<http://www.ncbi.nlm.nih.gov/pubmed/25338273>
- **Kim JG, Suyama S, Koch M, Jin S, Argente-Arizon P, Argente J, Liu ZW, Zimmer MR, Jeong JK, Szigeti-Buck K, Gao Y, Garcia-Caceres C, Yi CX, Salmaso N, Vaccarino FM, Chowen J, Diano S, Dietrich MO, Tschöp MH, Horvath TL.** Leptin signaling in astrocytes regulates hypothalamic neuronal circuits and feeding. *Nat Neurosci* 2014; 17(7):908-10
<http://www.ncbi.nlm.nih.gov/pubmed/24880214>
- **Kim JH, Meyers MS, Khuder MS, Abdallah SL, Muturi HT, Russo L, Tate CR, Hevener AL, Najjar SM, Leloup C, Mauvais-Jarvis F.** Tissue-selective estrogen complexes with bazedoxifene prevent metabolic dysfunction in female mice. *Mol Metab* 2014; 3(2):177-90
<http://www.ncbi.nlm.nih.gov/pubmed/24634829>
- **Kukat A, Dogan SA, Edgar D, Mourier A, Jacoby C, Maiti P, Mauer J, Becker C, Senft K, Wibom R, Kudin AP, Hultenby K, Flögel U, Rosenkranz S, Ricquier D, Kunz WS, Trifunovic A.** Loss of UCP2 attenuates mitochondrial dysfunction without altering ROS production and uncoupling activity. *PLoS Genet* 2014; 10(6):e1004385
<http://www.ncbi.nlm.nih.gov/pubmed/24945157>
- **Laeger T, Henagan TM, Albarado DC, Redman LM, Bray GA, Noland RC, Münzberg H, Hutson SM, Gettys TW, Schwartz MW, Morrison CD.** FGF21 is an endocrine signal of protein restriction. *J Clin Invest* 2014; 124(9):3913-22
<http://www.ncbi.nlm.nih.gov/pubmed/25133427>
- **Liu C, Bookout AL, Lee S, Sun K, Jia L, Lee C, Udit S, Deng Y, Scherer PE, Mangelsdorf DJ, Gautron L, Elmquist JK.** PPAR γ in vagal neurons regulates high-fat diet induced thermogenesis. *Cell Metab* 2014; 19(4):722-30
<http://www.ncbi.nlm.nih.gov/pubmed/24703703>
- **Mauer J, Chaurasia B, Goldau J, Vogt MC, Ruud J, Nguyen KD, Theurich S, Hausen AC, Schmitz J, Brönneke HS, Estevez E, Allen TL, Mesaros A, Partridge L, Febbraio MA, Chawla A, Wunderlich FT, Brüning JC.** Signaling by IL-6 promotes alternative activation of macrophages to limit endotoxemia and obesity-associated resistance to insulin. *Nat Immunol* 2014; 15(5):423-30
<http://www.ncbi.nlm.nih.gov/pubmed/24681566>
- **McAllan L, Skuse P, Cotter PD, O'Connor P, Cryan JF, Ross RP, Fitzgerald G, Roche HM, Nilaweera KN.** Protein quality and the protein to carbohydrate ratio within a high fat diet influences energy balance and the gut microbiota in C57BL/6J mice. *PLoS One* 2014; 9(2):e88904

<http://www.ncbi.nlm.nih.gov/pubmed/24520424>

- **Müller-Fielitz H, Hübel N, Mildner M, Vogt FM, Barkhausen J, Raasch W.** Chronic blockade of angiotensin AT1 receptors improves cardinal symptoms of metabolic syndrome in diet-induced obesity in rats. *Br J Pharmacol* 2014; 171(3):746-60
<http://www.ncbi.nlm.nih.gov/pubmed/24490862>
- **Ravussin Y, LeDuc CA, Watanabe K, Mueller BR, Skowronski A, Rosenbaum M, Leibel RL.** Effects of chronic leptin infusion on subsequent body weight and composition in mice: Can body weight set point be reset? *Mol Metab* 2014; 3(4):432-40
<http://www.ncbi.nlm.nih.gov/pubmed/24944902>
- **Soares FA, Dalla Corte CL, Andrade ER, Marina R, González P, Barrio JP.** Purple grape juice as a protector against acute x-irradiation induced alterations on mobility, anxiety, and feeding behaviour in mice. *Nutr Hosp* 2014; 29(4):812-21
<http://www.ncbi.nlm.nih.gov/pubmed/24679021>
- **Spieler D, Kaffe M, Knauf F, Bessa J, Tena JJ, Giesert F, Schormair B, Tilch E, Lee H, Horsch M, Czamara D, Karbalai N, von Toerne C, Waldenberger M, Gieger C, Lichtner P, Claussnitzer M, Naumann R, Müller-Myhsok B, Torres M, Garrett L, Rozman J, Klingenspor M, Gailus-Durner V, Fuchs H, Hrabě de Angelis M, Beckers J, Hölter SM, Meitinger T, Hauck SM, Laumen H, Wurst W, Casares F, Gómez-Skarmeta JL, Winkelmann J.** Restless legs syndrome-associated intronic common-variant in *Meis1* alters enhancer function in the developing telencephalon. *Genome Res* 2014; 24(4):592-603
<http://www.ncbi.nlm.nih.gov/pubmed/24642863>
- **Stoffel W, Hammels I, Jenke B, Binczek E, Schmidt-Soltau I, Brodesser S, Odenthal M, Thevis M.** Obesity resistance and deregulation of lipogenesis in D6-fatty acid desaturase (*FADS2*) deficiency. *EMBO Rep* 2014; 15(1):110-20
<http://www.ncbi.nlm.nih.gov/pubmed/24378641>
- **Suárez J, Rivera P, Arrabal S, Crespillo A, Serrano A, Baixeras E, Pavón FJ, Cifuentes M, Nogueiras R, Ballesteros J, Dieguez C, Rodríguez de Fonseca F.** Oleoylethanolamide enhances β -adrenergic-mediated thermogenesis and white-to-brown adipocyte phenotype in epididymal white adipose tissue in rat. *Dis Model Mech* 2014; 7(1):129-41
<http://www.ncbi.nlm.nih.gov/pubmed/24159189>
- **Sudakov SK, Nazarova GA, Alekseeva EV, Kolpakov AA.** Effect of activated peripheral κ -opioid receptors on the action of nicotine and its withdrawal in nicotine-dependent rats. *Bull Exp Biol Med* 2014; 156(5):609-11
<http://www.ncbi.nlm.nih.gov/pubmed/24770739>
- **Takeda Y, Kang HS, Freudenberg J, DeGraff LM, Jothi R, Jetten AM.** Retinoic acid-related orphan receptor γ (ROR γ): a novel participant in the diurnal regulation of hepatic gluconeogenesis and insulin sensitivity. *PLoS Gent* 2014; 10(5):e1004331
<http://www.ncbi.nlm.nih.gov/pubmed/24831725>
- **Torrens JM, Orellana-Gavaldà JM, Palou M, Sánchez J, Herrero L, Picó C, Serra D, Palou A.** Enhancing hepatic fatty acid oxidation as a strategy for reversing metabolic disorders programmed by maternal undernutrition during gestation. *Cell Physiol Biochem* 2014; 33(5):1498-515
<http://www.ncbi.nlm.nih.gov/pubmed/24854839>

- **Trigub MM, Bogdanova NG, Kolpakov AA, Bashkatova VG, Sudakov SK.** Effect of peripheral opioid receptor agonists on depressive activity of ethanol. *Bull Exp Biol Med* 2014; 156(6):778-80
<http://www.ncbi.nlm.nih.gov/pubmed/24824695>
- **Tsaousidou E, Paeger L, Belgardt BF, Pal M, Wunderlich CM, Brönneke H, Collienne U, Hampel B, Wunderlich FT, Schmidt-Supprian M, Kloppenburg P, Brüning JC.** Distinct Roles for JNK and IKK Activation in Agouti- Related Peptide Neurons in the Development of Obesity and Insulin Resistance. *Cell Rep* 2014; 9(4):1495-506
<http://www.ncbi.nlm.nih.gov/pubmed/25456138>
- **Tucci V, Kleefstra T, Hardy A, Heise I, Maggi S, Willemsen MH, Hilton H, Esapa C, Simon M, Buenavista MT, McGuffin LJ, Vizer L, Doderio L, Tsiftaris S, Romero R, Nillesen WN, Vissers LE, Kempers MJ, Vulto-van Silfhout AT, Iqbal Z, Orlando M, Maccione A, Lassi G, Farisello P, Constestabile A, Tinarelli F, Nieuw T, Raimondi A, Greco B, Cantatore D, Gasparini L, Berdondini L, Bifone A, Gozzi A, Wells S, Nolan PM.** Dominant β -catenin mutations cause intellectual disability with recognizable syndromic features. *J Clin Invest* 2014; 124(4):1468-82
<http://www.ncbi.nlm.nih.gov/pubmed/24614104>
- **Urbach YK, Raber KA, Canneva F, Plank AC, Andreasson T, Ponten H, Kullingsjö J, Nguyen HP, Riess O, von Hörsten S.** Automated phenotyping and advanced data mining exemplified in rats transgenic for Huntington's disease. *J Neurosci Methods* 2014; 234:38-53
<http://www.ncbi.nlm.nih.gov/pubmed/25020253>
- **Weimer S, Priebes J, Kuhlow D, Groth M, Priebe S, Mansfeld J, Merry TL, Dubuis S, Laube B, Pfeiffer AF, Schulz TJ, Guthke R, Platzer M, Zamboni N, Zarse K, Ristow M.** D-Glucosamine supplementation extends life span of nematodes and of ageing mice. *Nat Commun* 2014; 5:3563
<http://www.ncbi.nlm.nih.gov/pubmed/24714520>
- **Williams EG, Mouchiroud L, Frochoux M, Pandey A, Andreux PA, Deplancke B, Auwerx J.** An evolutionarily conserved role for the aryl hydrocarbon receptor in the regulation of movement. *PLoS Genet* 2014; 10(9):e1004673
<http://www.ncbi.nlm.nih.gov/pubmed/25255223>
- **Wuest S, Mueller R, Blüher M, Item F, Chin AS, Wiedemann MS, Takizawa H, Kovtonyuk L, Chervonsky AV, Schoenle EJ, Manz MG, Konrad D.** Fas (CD95) expression in myeloid cells promotes obesity-induced muscle insulin resistance. *EMBO Mol Med* 2014; 6(1):43-56
<http://www.ncbi.nlm.nih.gov/pubmed/24203314>

2013

- **Anthwal N, Pelling M, Claxton S, Mellitzer G, Collin C, Kessar N, Richardson WD, Gradwohl G, Ang SL.** Conditional deletion of neurogenin-3 using Nkx2.1iCre results in a mouse model for the central control of feeding, activity and obesity. *Dis Model Mech* 2013; 6(5):1133-45
<http://www.ncbi.nlm.nih.gov/pubmed/23649822>

- **Beiroa D, Romero-Picó A, Langa C, Bernabeu C, López M, López-Novoa JM, Nogueiras R, Diéguez C.** Heterozygous deficiency of endoglin decreases insulin and hepatic triglyceride levels during high fat diet. *PLoS one* 2013; 8(1): e54591
<http://www.ncbi.nlm.nih.gov/pubmed/23336009>
- **Berglund ED, Liu C, Sohn JW, Liu T, Kim MH, Lee CE, Vianna CR, Williams KW, Xu Y, Elmquist JK.** Serotonin 2C receptors in pro-opiomelanocortin neurons regulate energy and glucose homeostasis. *J Clin Invest* 2013; 123(12):5061-70
<http://www.ncbi.nlm.nih.gov/pubmed/24177424>
- **Binder E, Bermúdez-Silva FJ, André C, Elie M, Romero-Zerbo SY, Leste-Lasserre T, Belluomo L, Duchamp A, Clark S, Aubert A, Mezzullo M, Fanelli F, Pagotto U, Layé S, Mithieux G, Cota D.** Leucine supplementation protects from insulin resistance by regulating adiposity levels. *PLoS One* 2013; 8(9):e74705
<http://www.ncbi.nlm.nih.gov/pubmed/24086364>
- **Boon MR, van den Berg SA, Wang Y, van den Bossche J, Karkampouna S, Bauwens M, De Saint-Hubert M, van der Horst G, Vukicevic S, de Winther MP, Havekes LM, Jukema JW, Tamsma JT, van der Pluijm G, van Dijk KW, Rensen PC.** BMP7 activates brown adipose tissue and reduces diet-induced obesity only at subthermoneutrality. *PLoS One* 2013; 8(9):e74083
<http://www.ncbi.nlm.nih.gov/pubmed/24066098>
- **Dutia R, Kim AJ, Modes M, Rothlein R, Shen JM, Tian YE, Ihbais J, Victory SF, Valcarce C, Wardlaw SL.** Effects of AgRP inhibition on energy balance and metabolism in rodent models. *PLoS One* 2013; 8(6):e65317
<http://www.ncbi.nlm.nih.gov/pubmed/23762342>
- **Gao Y, Nelson DW, Banh T, Yen MI, Yen CL.** Intestine-specific expression of MOGAT2 partially restores metabolic efficiency in Mogat2-deficient mice. *J Lipid Res* 2013; 54(6): 1644-52
<http://www.ncbi.nlm.nih.gov/pubmed/23536640>
- **Greenman Y, Kuperman Y, Drori Y, Asa SL, Navon I, Forkosh O, Gil S, Stern N, Chen A.** Postnatal ablation of POMC neurons induces an obese phenotype characterized by decreased food intake and enhanced anxiety-like behavior. *Mol Endocrinol* 2013; 27(7):1091-102
<http://www.ncbi.nlm.nih.gov/pubmed/23676213>
- **Hanada T, Weitzer S, Mair B, Bernreuther C, Wainger BJ, Ichida J, Hanada R, Orthofer M, Cronin SJ, Komnenovic V, Minis A, Sato F, Mimata H, Yoshimura A, Tamir I, Rainer J, Kofler R, Yaron A, Eggan KC, Woolf CJ, Glatzel M, Herbst R, Martinez J, Penninger JM.** CLP1 links tRNA metabolism to progressive motor-neuron loss. *Nature* 2013; 495(7442):474-80
<http://www.ncbi.nlm.nih.gov/pubmed/23474986>
- **Harley IT, Giles DA, Pfluger PT, Burgess SL, Walters S, Hembree J, Raver C, Rewerts CL, Downey J, Flick LM, Stankiewicz TE, McAlees JW, Wills-Karp M, Balfour Sartor R, Divanovic S, Tschöp MH, Karp CL.** Differential colonization with segmented filamentous bacteria and *Lactobacillus murinus* do not drive divergent development of diet-induced obesity in C57BL/6 mice. *Mol Metab* 2013; 2(3):171-83
<http://www.ncbi.nlm.nih.gov/pubmed/24049732>

- **Heckmann BL, Zhang X, Xie X, Saarinen A, Lu X, Yang X, Liu J.** Defective adipose lipolysis and altered global energy metabolism in mice with adipose overexpression of the lipolytic inhibitor G0/G1 Switch Gene 2. *J Biol Chem* 2013; Epub <http://www.ncbi.nlm.nih.gov/pubmed/24302733>
- **Holst B, Madsen KL, Jansen AM, Jin C, Rickhag M, Lund VK, Jensen M, Bhatia V, Sørensen G, Madsen AN, Xue Z, Møller SK, Woldbye D, Qvortrup K, Hukanir R, Stamou D, Kjærulff O, Gether U.** PICK1 deficiency impairs secretory vesicle biogenesis and leads to growth retardation and decreased glucose tolerance. *PLoS One* 2013; 11(4):e1001542 <http://www.ncbi.nlm.nih.gov/pubmed/23630454>
- **Jiao J, Bae EJ, Bandyopadhyay G, Oliver J, Marathe C, Chen M, Hsu JY, Chen Y, Tian H, Olefsky JM, Saberi M.** Restoration of euglycemia after duodenal bypass surgery is reliant on central and peripheral inputs in Zucker fa/farats. *Diabetes* 2013; 62(4):1074-83 <http://www.ncbi.nlm.nih.gov/pubmed/23248171>
- **Jung DY, Chalasani U, Pan N, Friedline RH, Prosdocimo DA, Nam M, Azuma Y, Maganti R, Yu K, Velagapudi A, O'Sullivan-Murphy B, Sartoretto JL, Jain MK, Cooper MP, Urano F, Kim JK, Gray S.** KLF15 is a molecular link between endoplasmic reticulum stress and insulin resistance. *PLoS One* 2013; 8(10):e77851 <http://www.ncbi.nlm.nih.gov/pubmed/24167585>
- **Kant S, Barrett T, Vertii A, Noh YH, Jung DY, Kim JK, Davis RJ.** Role of the mixed-lineage protein kinase pathway in the metabolic stress response to obesity. *Cell Rep* 2013; 4(4):681-9 <http://www.ncbi.nlm.nih.gov/pubmed/23954791>
- **Kelp A, Koeppen AH, Petrasch-Parwez E, Calaminus C, Bauer C, Portal E, Yu-Taeger L, Pichler B, Bauer P, Riess O, Nguyen HP.** A Novel Transgenic Rat Model for Spinocerebellar Ataxia Type 17 Recapitulates Neuropathological Changes and Supplies In Vivo Imaging Biomarkers. *J Neurosci* 2013; 33(21):9068-81 <http://www.ncbi.nlm.nih.gov/pubmed/23699518>
- **Kirchner H, Heppner KM, Holland J, Kabra D, Tschöp MH, Pfluger PT.** Ablation of ghrelin O-acyltransferase does not improve glucose intolerance or body adiposity in mice on a leptin-deficient ob/ob background. *PLoS one* 2013; 8(4):e61822 <http://www.ncbi.nlm.nih.gov/pubmed/23630616>
- **Krüger J, Trappiel M, Dagnell M, Stawowy P, Meyborg H, Böhm C, Bhanot S, Ostman A, Kintscher U, Kappert K.** Targeting density-enhanced phosphatase-1 (DEP-1) with antisense oligonucleotides improves the metabolic phenotype in high-fat diet-fed mice. *Cell Commun Signal* 2013; 11(1):49 <http://www.ncbi.nlm.nih.gov/pubmed/23889985>
- **Mattjssen F, Alex S, Swarts HJ, Groen AK, van Schothorst EM, Kersten S.** Angptl4 serves as an endogenous inhibitor of intestinal lipid digestion. *Molecular Metabolism* 2013; 3(2):135-44 <http://www.ncbi.nlm.nih.gov/pubmed/24634819>

- **Müller TD, Lee SJ, Jastroch M, Kabra D, Stemmer K, Aichler M, Abplanalp B, Ananthakrishnan G, Bhardwaj N, Collins S, Divanovic S, Ende M, Finan B, Gao Y, Habegger KM, Hembree J, Heppner KM, Hofmann S, Holland J, Kuchler D, Kutschke M, Krishna R, Lehti M, Oelkrug R, Ottaway N, Perez-Tilve D, Raver C, Walch AK, Schriever SC, Speakman J, Tseng YH, Diaz-Meco M, Pfluger PT, Moscat J, Tschöp MH.** p62 links β -adrenergic input to mitochondrial function and thermogenesis. *The Journal of clinical investigation* 2013; 123(1): 469-78
<http://www.ncbi.nlm.nih.gov/pubmed/23257354>
- **Painsipp E, Köfer MJ, Farzi A, Dischinger US, Sinner F, Herzog H, Holzer P.** Neuropeptide Y and peptide YY protect from weight loss caused by Bacille Calmette-Guérin in mice. *Br J Pharmacol* 2013; 170(5):1014-26
<http://www.ncbi.nlm.nih.gov/pubmed/23992467>
- **Pal M, Wunderlich CM, Spohn G, Brönneke HS, Schmidt-Supprian M, Wunderlich FT.** Alteration of JNK-1 signaling in skeletal muscle fails to affect glucose homeostasis and obesity-associated insulin resistance in mice. *PloS one* 2013; 8(1): e54247
<http://www.ncbi.nlm.nih.gov/pubmed/23349837>
- **Pazos P, Lima L, Casanueva FF, Diéguez C, García MC.** Interleucine 6 deficiency modulates the hypothalamic expression of energy balance regulating peptides during pregnancy in mice. *PLoS One* 2013; 8(8):e72339
<http://www.ncbi.nlm.nih.gov/pubmed/24015235>
- **Perez-Sieira S, Martinez G, Porteiro B, Lopez M, Vidal A, Nogueiras R, Dieguez C.** Female Nur77-deficient mice show increased susceptibility to diet-induced obesity. *PloS one* 2013; 8(1): e53836
<http://www.ncbi.nlm.nih.gov/pubmed/23342015>
- **Picard A, Rouch C, Kassis N, Moulle VS, Croizier S, Denis RG, Castel J, Coant N, Davis K, Clegg DJ, Benoit SC, Prevot V, Bouret S, Luquet S, Le Stunff H, Cruciani-Guglielmacci C, Magnan C.** Hippocampal lipoprotein lipase regulates energy balance in rodents. *Molecular Metabolism* 2013; 3(2):167-76
<http://www.ncbi.nlm.nih.gov/pubmed/24634821>
- **Ravnskjaer K, Hogan MF, Lackey D, Tora L, Dent SY, Olefsky J, Montminy M.** Glucagon regulates gluconeogenesis through KAT2B- and WDR5-mediated epigenetic effects. *J Clin Invest* 2013; 123(19):4318-28
<http://www.ncbi.nlm.nih.gov/pubmed/24051374>
- **Ravussin Y, Gutman R, LeDuc CA, Leibel RL.** Estimating energy expenditure in mice using an energy balance technique. *International Journal of Obesity* 2013; 37(3): 399-403
<http://www.ncbi.nlm.nih.gov/pubmed/22751256>
- **Ren H, Plum-Morschel L, Gutierrez-Juarez R, Lu TY, Kim-Muller JY, Heinrich G, Wardlaw SL, Silver R, Accili D.** Blunted refeeding response and increased locomotor activity in mice lacking FoxO1 in synapsin-cre-expressing neurons. *Diabetes* 2013; 62(10):3373-83
<http://www.ncbi.nlm.nih.gov/pubmed/23835335>
- **Robinson L, Plano A, Cobb S, Riedel G.** Long-term home cage activity scans reveal lowered exploratory behaviour in symptomatic female Rett mice. *Behav Brain Res* 2013; 250:148-56
<http://www.ncbi.nlm.nih.gov/pubmed/23643691>

- **Rouger L, Denis GR, Luangsay S, Parmentier M.** Chem R23 knockout mice display mild obesity but no deficit in adipocyte differentiation. *J Endocrinol* 2013; 2019(3):279-89
<http://www.ncbi.nlm.nih.gov/pubmed/24084834>
- **Shechter R, London A, Kuperman Y, Ronen A, Rolls A, Chen A, Schwartz M.** Hypothalamic neuronal toll-like receptor 2 protects against age-induced obesity. *Sci Rep* 2013; 3:1254
<http://www.ncbi.nlm.nih.gov/pubmed/23409245>
- **Sims EK, Hatanaka M, Morris DL, Tersey SA, Kono T, Chaudry ZZ, Day KH, Moss DR, Stull ND, Mirmira RG, Evans-Molina C.** Divergent compensatory responses to high-fat diet between C57BL6/J and C57BLKS/J inbred mouse strains. *Am J Physiol Endocrinol Metab* 2013; 305(12): E1495-511
<http://www.ncbi.nlm.nih.gov/pubmed/24169046>
- **Sudakov SK, Bashkatova VG.** Effect of peripheral D₂ dopamine receptor antagonist domperidone on metabolism, feeding behavior, and locomotor activity of rats. *Bull Exp Biol Med* 2013; 155(6):705-7
<http://www.ncbi.nlm.nih.gov/pubmed/24288745>
- **Sun K, Halberg N, Khan M, Magalang UJ, Scherer PE.** Selective inhibition of hypoxia-inducible factor 1 α ameliorates adipose tissue dysfunction. *Mol Cell Biol* 2013; 33(5):904-17
<http://www.ncbi.nlm.nih.gov/pubmed/23249949>
- **Tang G-B, Tang X-F, Li K, Wang D-H.** Role of leptin in metabolic adaptation during cold acclimation. *J Stress Physiol Biochem* 2013; 9(4):101-14
http://www.jspb.ru/issues/2013/N4/JSPB_2013_4_101-114.pdf
- **Varcoe TJ, Boden MJ, Voultzios A, Salkeld MD, Rattanatray L, Kennaway DJ.** Characterisation of the maternal response to chronic phase shifts during gestation in the rat: implications for fetal metabolic programming. *PLoS One* 2013; 8(1):e53800
<http://www.ncbi.nlm.nih.gov/pubmed/23342007>
- **Veyrat-Durebex C, Poher AL, Caillon A, Somm E, Vallet P, Charnay Y, Rohner-Jeanrenaud F.** Improved Leptin Sensitivity as a Potential Candidate Responsible for the Spontaneous Food Restriction of the Lou/C Rat. *PLoS One* 2013; 8(9):e73452
<http://www.ncbi.nlm.nih.gov/pubmed/24039946>

- **Watanabe K, Watson E, Cremona ML, Millings EJ, Lefkowitz JH, Fischer SG, LeDuc CA, Leibel RL.** ILDR2: an endoplasmic reticulum resident molecule mediating hepatic lipid homeostasis. *PLoS One* 2013; 6(8):e67234
<http://www.ncbi.nlm.nih.gov/pubmed/23826244>
- **Wuest S, Mueller R, Blüher M, Item F, Chin AS, Wiedemann MS, Takizawa H, Kovtonyuk L, Chervonsky AV, Schoenle EJ, Manz MG, Konrad D.** Fas (CD95) expression in myeloid cells promotes obesity-induced muscle insulin resistance. *EMBO Mol Med* 2013; Epub
<http://www.ncbi.nlm.nih.gov/pubmed/24203314>
- **Yo K, Yu YM, Zhao G, Bonab AA, Aikawa N, Tompkins RG, Fischman AJ.** Brown adipose tissue and its modulation by a mitochondria-targeted peptide in rat burn injury-induced hypermetabolism. *Am J Physiol Endocrinol Metab* 2013; 304(4):E331-41
<http://www.ncbi.nlm.nih.gov/pubmed/23169784>
- **Zierler KA, Jaeger D, Pollak NM, Eder S, Rechberger GN, Radner FP, Woelkart G, Kolb D, Schmidt A, Kumari M, Preiss-Landl K, Pieske B, Mayer B, Zimmermann R, Lass A, Zechner R, Haemmerle G.** Functional cardiac lipolysis in mice critically depends on comparative gene identification-58. *J Biol Chem* 2013; 288(14):9892-904
<http://www.ncbi.nlm.nih.gov/pubmed/23413028>

2012

- **Batista TM, Alonso-Magdalena P, Vieira E, Amaral ME, Cederroth CR, Nef S, Quesada I, Carneiro EM, Nadal A.** Short-term treatment with bisphenol-A leads to metabolic abnormalities in adult male mice. *PloS one* 2012; 7(3): e33814
<http://www.ncbi.nlm.nih.gov/pubmed/22470480>
- **Benz V, Bloch M, Wardat S, Böhm C, Maurer L, Mahmoodzadeh S, Wiedmer P, Spranger J, Foryst-Ludwig A, Kintscher U.** Sexual dimorphic regulation of body weight dynamics and adipose tissue lipolysis. *PloS one* 2012; 7(5): e37794
<http://www.ncbi.nlm.nih.gov/pubmed/22662224>
- **Betz MJ, Bielohuby M, Mauracher B, Abplanalp W, Müller HH, Pieper K, Ramisch J, Tschöp MH, Beuschlein F, Bidlingmaier M, Slawik M.** Isoenergetic feeding of low carbohydrate-high fat diets does not increase brown adipose tissue thermogenic capacity in rats. *PloS one* 2012; 7(6): e38997
<http://www.ncbi.nlm.nih.gov/pubmed/22720011>
- **Blum ID, Lamont EW, Rodrigues T, Abizaid A.** Isolating neural correlates of the pacemaker for food anticipation. *PloS one* 2012; 7(4): e36117
<http://www.ncbi.nlm.nih.gov/pubmed/22558352>
- **Cai L, Wang Z, Ji A, Meyer JM, van der Westhuyzen DR.** Scavenger receptor CD36 expression contributes to adipose tissue inflammation and cell death in diet-induced obesity. *PloS one* 2012; 7(5): e36785
<http://www.ncbi.nlm.nih.gov/pubmed/22615812>

- **Castañeda TR, Abplanalp W, Um SH, Pfluger PT, Schrott B, Brown K, Grant E, Carnevalli L, Benoit SC, Morgan DA, Gilham D, Hui DY, Rahmouni K, Thomas G, Kozma SC, Clegg DJ, Tschöp MH.** Metabolic control by S6 kinases depends on dietary lipids. *PloS one* 2012; 7(3): e32631
<http://www.ncbi.nlm.nih.gov/pubmed/22412899>
- **Deblon N, Bourgoïn L, Veyrat-Durebex C, Peyrou M, Vinciguerra M, Caillon A, Maeder C, Fournier M, Montet X, Rohner-Jeanrenaud F, Foti M.** Chronic mTOR inhibition by rapamycin induces muscle insulin resistance despite weight loss in rats. *British journal of pharmacology* 2012; 165(7): 2325-40
<http://www.ncbi.nlm.nih.gov/pubmed/22014210>
- **Dietrich MO, Bober J, Ferreira JG, Tellez LA, Mineur YS, Souza DO, Gao XB, Picciotto MR, Araújo I, Liu ZW, Horvath TL.** AgRP neurons regulate development of dopamine neuronal plasticity and nonfood-associated behaviors. *Nature neuroscience* 2012; 15(8): 1108-10
<http://www.ncbi.nlm.nih.gov/pubmed/22729177>
- **Finley LW, Lee J, Souza A, Desquirit-Dumas V, Bullock K, Rowe GC, Procaccio V, Clish CB, Arany Z, Haigis MC.** Skeletal muscle transcriptional coactivator PGC-1 α mediates mitochondrial, but not metabolic, changes during calorie restriction. *Proceedings of the National Academy of Sciences of the United States of America* 2012; 109(8): 2931-6
<http://www.ncbi.nlm.nih.gov/pubmed/22308395>
- **Fournier B, Murray B, Gutzwiller S, Marcaletti S, Marcellin D, Bergling S, Brachat S, Persohn E, Pierrel E, Bombard F, Hatakeyama S, Trendelenburg AU, Morvan F, Richardson B, Glass DJ, Lach-Trifilieff E, Feige JN.** Blockade of the activin receptor IIb activates functional brown adipogenesis and thermogenesis by inducing mitochondrial oxidative metabolism. *Molecular and cellular biology* 2012; 32(14): 2871-9
<http://www.ncbi.nlm.nih.gov/pubmed/22586266>
- **Habegger KM, Matzke D, Ottaway N, Hembree J, Holland J, Raver C, Mansfeld J, Müller TD, Perez-Tilve D, Pfluger PT, Lee SJ, Diaz-Meco M, Moscat J, Leitges M, Tschöp MH, Hofmann SM.** Role of adipose and hepatic atypical protein kinase C lambda (PKC λ) in the development of obesity and glucose intolerance. *Adipocyte* 2012; 1(4): 203-214
<http://www.ncbi.nlm.nih.gov/pubmed/23700535>
- **Hassouna R, Zizzari P, Viltart O, Yang SK, Gardette R, Videau C, Badoer E, Epelbaum J, Tolle V.** A natural variant of obestatin, Q90L, inhibits ghrelin's action on food intake and GH secretion and targets NPY and GHRH neurons in mice. *PloS one* 2012; 7(12): e51135
<http://www.ncbi.nlm.nih.gov/pubmed/23251435>
- **He C, Bassik MC, Moresi V, Sun K, Wei Y, Zou Z, An Z, Loh J, Fisher J, Sun Q, Korsmeyer S, Packer M, May HI, Hill JA, Virgin HW, Gilpin C, Xiao G, Bassel-Duby R, Scherer PE, Levine B.** Exercise-induced BCL2-regulated autophagy is required for muscle glucose homeostasis. *Nature* 2012; 481(7382): 511-5
<http://www.ncbi.nlm.nih.gov/pubmed/22258505>
- **Hesse D, Jaschke A, Kanzleiter T, Witte N, Augustin R, Hommel A, Püschel GP, Petzke KJ, Joost HG, Schupp M, Schürmann A.** GTPase ARFRP1 is essential for normal hepatic glycogen storage and insulin-like growth factor 1 secretion. *Mol Cell Biol* 2012; 32(21):4363-74
<http://www.ncbi.nlm.nih.gov/pubmed/22927645>

- **Jeong JK, Szabo G, Raso GM, Meli R, Diano S.** Deletion of prolyl carboxypeptidase attenuates the metabolic effects of diet-induced obesity. *American Journal of Physiology, Endocrinology and Metabolism* 2012; 302(12): e1502-10
<http://www.ncbi.nlm.nih.gov/pubmed/22454290>
- **Jiao Y, George SK, Zhao Q, Hulver MW, Hutson SM, Bishop CE, Lu B.** Mex3c mutation reduces adiposity and increases energy expenditure. *Molecular and cellular biology* 2012; 32(21): 4350-62
<http://www.ncbi.nlm.nih.gov/pubmed/22927639>
- **Karavia EA, Papachristou DJ, Liopeta K, Triantafyllidou I, Dimitrakopoulos O, Kypreos KE.** Apolipoprotein A-I modulates Processes Associated with Diet-Induced Nonalcoholic Fatty Liver Disease in Mice. *Molecular Medicine* 2012; 18: 901-12
<http://www.ncbi.nlm.nih.gov/pubmed/22576368>
- **Kim KW, Donato J Jr, Berglund ED, Choi YH, Kohno D, Elias CF, Depinho RA, Elmquist JK.** FOXO1 in the ventromedial hypothalamus regulates energy balance. *The Journal of clinical investigation* 2012; 122(7): 2578-89
<http://www.ncbi.nlm.nih.gov/pubmed/22653058>
- **Kim SJ, Nian C, Karunakaran S, Clee SM, Isaacs CM, McIntosh CH.** GIP-Overexpressing Mice Demonstrate Reduced Diet-Induced Obesity and Steatosis, and Improved Glucose Homeostasis. *Public Library of Science One* 2012; 7(7): e40156
<http://www.ncbi.nlm.nih.gov/pubmed/22802954>
- **Lassi G, Ball ST, Maggi S, Colonna G, Nieuw T, Cero C, Bartolomucci A, Peters J, Tucci V.** Loss of Gnas imprinting differentially affects REM/NREM sleep and cognition in mice. *PLoS Genet* 2012; 8(5): e1002706
<http://www.ncbi.nlm.nih.gov/pubmed/22589743>
- **Lee AW, Hengstler H, Schwald K, Berriel-Diaz M, Loreth D, Kirsch M, Kretz O, Haas CA, de Angelis MH, Herzig S, Brümmendorf T, Klingenspor M, Rathjen FG, Rozman J, Nicholson G, Cox RD, Schäfer MK.** Functional inactivation of the genome-wide association study obesity gene neuronal growth regulator 1 in mice causes a body mass phenotype. *PloS one* 2012; 7(7): e41537
<http://www.ncbi.nlm.nih.gov/pubmed/22844493>
- **Li Y, Mouche S, Sajic T, Veyrat-Durebex C, Supale R, Pierroz D, Ferrari S, Negro F, Hasler U, Feraille E, Moll S, Meda P, Deffert C, Montet X, Krause KH, Szanto I.** Deficiency in the NADPH oxidase 4 predisposes towards diet-induced obesity. *International Journal of Obesity* 2012; 36(12): 1503-13
<http://www.ncbi.nlm.nih.gov/pubmed/22430302>
- **Lockie SH, Heppner KM, Chaudhary N, Chabenne JR, Morgan DA, Veyrat-Durebex C, Ananthakrishnan G, Rohner-Jeanrenaud F, Drucker DJ, DiMarchi R, Rahmouni K, Oldfield BJ, Tschöp MH, Perez-Tilve D.** Direct control of brown adipose tissue thermogenesis by central nervous system glucagon-like peptide-1 receptor signaling. *Diabetes* 2012; 61(11): 2753-62
<http://www.ncbi.nlm.nih.gov/pubmed/22933116>
- **Mang GM, Dürst T, Bürki H, Imobersteg S, Abramowski D, Schuepbach E, Hoyer D, Fendt M, Gee CE.** The Dual Orexin Receptor Antagonist Almorexant Induces Sleep and Decreases Orexin-Induced Locomotion by Blocking Orexin 2 Receptors. *Sleep* 2012; 35(12): 1625-35

- <http://www.ncbi.nlm.nih.gov/pubmed/23204605>
- **Martínez de Morentin PB, Whittle AJ, Fernø J, Nogueiras R, Diéguez C, Vidal-Puig A, López M.** Nicotine induces negative energy balance through hypothalamic AMP-activated protein kinase. *Diabetes* 2012; 61(4): 807-17
<http://www.ncbi.nlm.nih.gov/pubmed/22315316>
 - **Mosienko V, Bert B, Beis D, Matthes S, Fink H, Bader M, Alenina N.** Exaggerated aggression and decreased anxiety in mice deficient in brain serotonin. *Translational psychiatry* 2012; 2: e122
<http://www.ncbi.nlm.nih.gov/pubmed/22832966>
 - **Muñoz R, Carmody JS, Stylopoulos N, Davis P, Kaplan LM.** Isolated duodenal exclusion increases energy expenditure and improves glucose homeostasis in diet-induced obese rats. *Am J Physiol Regul Integr Comp Physiol* 2012; 303(10): R985-93
<http://www.ncbi.nlm.nih.gov/pubmed/22972837>
 - **Olszewski PK, Rozman J, Jacobsson JA, Rathkolb B, Strömberg S, Hans W, Klockars A, Alsiö J, Risérus U, Becker L, Hölter SM, Elvert R, Ehrhardt N, Gailus-Durner V, Fuchs H, Fredriksson R, Wolf E, Klopstock T, Wurst W, Levine AS, Marcus C, de Angelis MH, Klingenspor M, Schiöth HB, Kilimann MW.** Neurobeachin, a regulator of synaptic protein targeting, is associated with body fat mass and feeding behavior in mice and body-mass index in humans. *PLoS genetics* 2012; 8(3): e1002568
<http://www.ncbi.nlm.nih.gov/pubmed/22438821>
 - **Pacheco-López G, Giovanoli S, Langhans W, Meyer U.** Priming of Metabolic Dysfunctions by Prenatal Immune Activation in Mice: Relevance to Schizophrenia. *Schizophrenia Bulletin* 2012; 39(2): 319-29
<http://www.ncbi.nlm.nih.gov/pubmed/22186136>
 - **Solverson P, Murali SG, Brinkman AS, Nelson DW, Clayton MK, Yen CL, Ney DM.** Glycomacropeptide, a low-phenylalanine protein isolated from cheese whey, supports growth and attenuates metabolic stress in the murine model of phenylketonuria. *Am J Physiol Endocrinol Metab* 2012; 302(7):e885-95
<http://www.ncbi.nlm.nih.gov/pubmed/22297302>
 - **Stefanidis A, Spencer SJ.** Effects of neonatal overfeeding on juvenile and adult feeding and energy expenditure in the rat. *PloS One* 2012; 7(12): e52130
<http://www.ncbi.nlm.nih.gov/pubmed/23251693>
 - **Van Klinken JB, van den Berg SA, Havekes LM, Willems Van Dijk K.** Estimation of activity related energy expenditure and resting metabolic rate in freely moving mice from indirect calorimetry data. *PloS One* 2012; 7(5): e36162
<http://www.ncbi.nlm.nih.gov/pubmed/22574139>
 - **Vianna CR, Donato J Jr, Rossi J, Scott M, Economides K, Gautron L, Pierpont S, Elias CF, Elmquist JK.** Cannabinoid receptor 1 in the vagus nerve is dispensable for body weight homeostasis but required for normal gastrointestinal motility. *J Neurosci* 2012; 32(30): 10331-7
<http://www.ncbi.nlm.nih.gov/pubmed/22836266>

2011

- **Al-Ahad Sabbagha NGA, Kao HJ, Yang CF, Huang CC, Lin WD, Tsai F, Chen TH, Tarn WY, Wu JY, Chen YT.** Alternative splicing in acad8 resulting a mitochondrial defect and progressive hepatic steatosis in mice. *Pediatric Research* 2011; 70(1): 31-6
http://www.scirp.org/journal/PaperInformation.aspx?paperID=50993#.VHLuhPmG_w8
- **Alexandru A, Jagla W, Graubner S, Becker A, Bäuscher C, Kohlmann S, Sedlmeier R, Raber KA, Cynis H, Rönicke R, Reymann KG, Petrasch-Parwez E, Hartlage-Rübsamen M, Waniek A, Rossner S, Schilling S, Osmand AP, Demuth HU, von Hörsten S.** Selective hippocampal neurodegeneration in transgenic mice expressing small amounts of truncated Abeta is induced by pyroglutamate-Abeta formation. *J Neurosci* 2011; 31(36): 12790-801
<http://www.ncbi.nlm.nih.gov/pubmed/21900558>
- **Boersma GJ, Benthem L, van Dijk G, Scheurink AJ.** Individual variation in the (patho)physiology of energy balance. *Physiol Behav* 2011; 103: 89-97
<http://www.ncbi.nlm.nih.gov/pubmed/21237186>
- **Deblon N, Veyrat-Durebex C, Bourgoin L, Caillon A, Bussier AL, Petrosino S, Piscitelli F, Legros JJ, Geenen V, Foti M, Wahli W, Di Marzo V, Rohner-Jeanrenaud F.** Mechanisms of the anti-obesity effects of oxytocin in diet-induced obese rats. *PLoS One* 2011; 6(9): e25565
<http://www.ncbi.nlm.nih.gov/pubmed/21980491>
- **Ferrero DM, Lemon JK, Fluegge D, Pashkovski SL, Korzan WJ, Datta SR, Spehr M, Fendt M, Liberles S.** Detection and avoidance of a carnivore odor by prey. *Proc Natl Acad Sci* 2011; 108(27): 11235-40
<http://www.ncbi.nlm.nih.gov/pubmed/21690383>
- **Haemmerle G, Moustafa T, Woelkart G, Büttner S, Schmidt A, van de Weijer T, Hesselink M, Jaeger D, Kienesberger PC, Zierler K, Schreiber R, Eichmann T, Kolb D, Kotzbeck P, Schweiger M, Kumari M, Eder S, Schoiswohl G, Wongsiriroj N, Pollak NM, Radner FPW, Preiss-Landl K, Kolbe T, Rüllicke T, Pieske B, Trauner M, Lass A, Zimmermann R, Hoefler G, Cinti S, Kershaw EE, Schrauwen P, Madeo F, Mayer B, Zechner R.** ATGL-mediated fat catabolism regulates cardiac mitochondrial function via PPAR-alpha and PGC-1. *Nat Med* 2011; 17(9): 1076-85
<http://www.ncbi.nlm.nih.gov/pubmed/21857651>
- **Hollis JH, Jonaidi H, Lemus M, Oldfield BJ.** The endocannabinoid Arachidonylethanolamide attenuates aspects of lipopolysaccharide-induced changes in energy intake, energy expenditure and hypothalamic Fos expression. *J Neuroimmunol* 2011; 233: 127-34
<http://www.ncbi.nlm.nih.gov/pubmed/21262543>
- **Hübener J, Vauti F, Funke C, Wolburg H, Ye Y, Schmidt T, Wolburg-Buchholz K, Schmitt I, Gardyan A, Drießen S, Arnold HH, Nguyen HP, Riess O.** N-terminal ataxin-3 causes neurological symptoms with inclusions, endoplasmic reticulum stress and ribosomal dislocation. *Brain* 2011; 134: 1925-42
<http://www.ncbi.nlm.nih.gov/pubmed/21653538>
- **Jamieson PM, Cleasby ME, Kuperman Y, Morton NM, Kelly PAT, Brownstein DG, Mustard KJ, Vaughan JM, Carter RN, Hahn CN, Hardie DG, Seckl JR, Chen A, Vale WW.** Urocortin 3 transgenic mice exhibit a metabolically favourable phenotype resisting obesity and hyperglycaemia on a high-fat diet. *Diabetologia* 2011; 54: 2392-403

<http://www.ncbi.nlm.nih.gov/pubmed/21667214>

- **Johnson LA, Arbones-Mainar JM, Fox RG, Pendse AA, Altenburg MK, Kim HS, Maeda N.** Apolipoprotein E4 exaggerates diabetic dyslipidemia and atherosclerosis in mice lacking the LDL receptor. *Diabetes* 2011; 60(9): 2285-94
<http://www.ncbi.nlm.nih.gov/pubmed/21810592>
- **Jordan SD, Krüger M, Willmes DM, Redemann N, Wunderlich FT, Brönneke HS, Merkwirth C, Kashkar H, Olkkonen VM, Böttger T, Braun T, Seibler J, Brüning JC.** Obesity-induced overexpression of miRNA-143 inhibits insulin-stimulated AKT activation and impairs glucose metabolism. *Nat Cell Biol* 2011; 13(4): 434-446
<http://www.ncbi.nlm.nih.gov/pubmed/21441927>
- **Levi J, Gray SL, Speck M, Huynh FK, Babich SL, Gibson WT, Kieffer TJ.** Acute disruption of leptin signaling in vivo leads to increased insulin levels and insulin resistance. *Endocrinology* 2011; 152(9): 3385-95
<http://www.ncbi.nlm.nih.gov/pubmed/21750049>
- **Li Y, Tong X, Rumala C, Clemons K, Wang S.** Thrombospondin1 deficiency reduces obesity-associated inflammation and improves insulin sensitivity in a diet-induced obese mouse model. *PloS One* 2011; 6(10): e26656
<http://www.ncbi.nlm.nih.gov/pubmed/22039525>
- **Lin HV, Ren H, Samuel VT, Lee HY, Lu TY, Shulman GI, Accili D.** Diabetes in mice with selective impairment of insulin action in Glut4-expressing tissues. *Diabetes* 2011; 60, 3: 700-708
<http://www.ncbi.nlm.nih.gov/pubmed/21266328>
- **Lutz SZ, Hennige AM, Feil S, Peter A, Gerling A, Machann J, Kröber SM, Rath M, Schürmann A, Weigert C, Häring HU, Feil R.** Genetic ablation of cGMP-dependent protein kinase type I causes liver inflammation and fasting hyperglycemia. *Diabetes* 2011; 60(5): 1566-76
<http://www.ncbi.nlm.nih.gov/pubmed/21464444>
- **Ma T, Liaset B, Hao Q, Koefoed Petersen R, Fjære E, Thi Ngo H, Haukûs Lillefosse H, Ringholm S, Brask Sonne S, Thue Treebak J, Pilegaard H, Frøyland L, Kristiansen K, Madsen L.** Sucrose counteracts the anti-inflammatory effect of fish oil in adipose tissue and increases obesity development in mice. *PloS One* 2011; 6(6): e21647
<http://www.ncbi.nlm.nih.gov/pubmed/21738749>
- **Nelson DW, Gao Y, Spencer NM, Banh T, Yen CLE.** Deficiency of MGAT2 increases energy expenditure without high-fat feeding and protects genetically obese mice from excessive weight gain. *J Lipid Res* 2011; 52: 1723-32
<http://www.ncbi.nlm.nih.gov/pubmed/21734185>
- **Nuber S, Petrasch-Parwez E, Arias-Carrión O, Koch L, Kohl Z, Schneider J, Calaminus C, Dermietzel R, Samarina A, Boy J, Nguyen HP, Teisman P, Palanichamy Velavan T, Kahle PJ, von Hörsten S, Fendt M, Krüger R, Riess O.** Olfactory neuron-specific expression of A30P alpha-synuclein exacerbates dopamine deficiency and hyperactivity in a novel conditional model of early parkinson's disease stages. *Neurobiol Dis* 2011; 44: 192-204
<http://www.ncbi.nlm.nih.gov/pubmed/21767644>

- **Pfluger PT, Castañeda TR, Heppner KM, Strassburg S, Kruthaupt T, Chaudhary N, Halem H, Culler MD, Datta R, Burget L, Tschöp MH, Nogueiras R, Perez-Tilve D.** Ghrelin, peptide YY and their hypothalamic targets differentially regulate spontaneous physical activity. *Physiol Behav* 2011; 105: 52-61
<http://www.ncbi.nlm.nih.gov/pubmed/21554896>
- **Ravussin Y, Gutman R, Diano S, Shanabrough M, Borok E, Sarman B, Lehmann A, LeDuc CA, Rosenbaum M, Horvarth TL, Leibel RL.** Effects of chronic weight perturbation on energy homeostasis and brain structure in mice. *Am J Physiol - Regul Integr Comp Physiol* 2011; 300: 1352-62
<http://www.ncbi.nlm.nih.gov/pubmed/21411766>
- **Rossi J, Balthasar N, Olson D, Scott M, Berglund E, Lee CE, Choi MJ, Lauzon D, Lowell BB, Elmquist JK.** Melanocortin-4 receptors expressed by cholinergic neurons regulate energy balance and glucose homeostasis. *Cell Metab* 2011; 13: 195-204
<http://www.ncbi.nlm.nih.gov/pubmed/21284986>
- **Rousso-Noori L, Knobler H, Levy-Apter E, Kuperman Y, Neufeld-Cohen A, Keshet Y, Akepati VR, Klinghoffer RA, Chen A, Elson A.** Protein tyrosine phosphatase epsilon affects body weight by downregulating leptin signaling in a phosphorylation-dependent manner. *Cell Metab* 2011; 13: 562-572
<http://www.ncbi.nlm.nih.gov/pubmed/21531338>
- **Savic D, Ye H, Aneas I, Park SY, Bell GI, Nobrega MA.** Alterations in TCF7L2 expression define its role as a key regulator of glucose metabolism. *Genome Res* 2011; 21(9): 1417-25
<http://www.ncbi.nlm.nih.gov/pubmed/21673050>
- **Schilling S, Kohlmann S, Bäuscher C, Sedlmeier R, Koch B, Eichentopf R, Becker A, Cynis H, Hoffmann T, Berg S, Freyse J, von Hörsten S, Rossner S, Graubner S, Demuth HU.** Glutaminy cyclase (OC) knock out mice show mild hypothyroidism but absence of hypogonadism: implications for enzyme function and drug development. *J Biol Chem* 2011; 286(16): 1-19
<http://www.ncbi.nlm.nih.gov/pubmed/21330373>
- **Schulz N, Himmelbauer H, Rath M, van Weeghel M, Houten S, Kulik W, Suhre K, Scherneck S, Vogel H, Kluge R, Wiedmer P, Joost HG, Schürmann A.** Role of Medium- and Short-Chain L-3-Hydroxyacyl- CoA Dehydrogenase in the Regulation of Body Weight and Thermogenesis. *Endocrinol* 2011; 152(12):4641-51
<http://www.ncbi.nlm.nih.gov/pubmed/21990309>
- **Scott MM, Williams KW, Rossi J, Lee CE, Elmquist JK.** Leptin receptor expression in hindbrain Glp-1 neurons regulates food intake and energy balance in mice. *J Clin Invest* 2011; 121(6): 2413-21
<http://www.ncbi.nlm.nih.gov/pubmed/21606595>
- **Petersen PS, Jin C, Nygaard Madsen A, Rasmussen M, Kuhre R, Egerod KL, Bo Nielsen L, Schwartz TW, Holst B.** Deficiency of the GPR39 receptor is associated with obesity and altered adipocyte metabolism. *FASEB J* 2011; 25: 1-12
<http://www.ncbi.nlm.nih.gov/pubmed/21784784>

- **Taschler U, Radner FPW, Heier C, Schreiber R, Schweiger M, Schoiswohl G, Preiss-Landl K, Jaeger D, Reiter B, Koefeler HC, Wojciechowski J, Theussl C, Penninger JM, Lass A, Haemmerle G, Rechner R, Zimmermann R.** Monoglyceride lipase deficiency in mice impairs lipolysis and attenuates diet-induced insulin resistance. *J Biol Chem* 2011; 286(20): 17467-77
<http://www.ncbi.nlm.nih.gov/pubmed/21454566>
- **Tauriainen E, Luostarinen M, Martonen E, Finckenberg P, Kovalainen M, Huotari A, Herzig KH, Lecklin A, Mervaala E.** Distinct effects of calorie restriction and resveratrol on diet-induced obesity and Fatty liver formation. *J Nutr Metab* 2011; 2011: 525094
<http://www.ncbi.nlm.nih.gov/pubmed/21977315>
- **Tschöp MH, Speakman JR, Arch JR, Auwerx J, Brüning JC, Chan L, Eckel RH, Farese RV Jr, Galgani JE, Hambly C, Herman MA, Horvath TL, Kahn BB, Kozma SC, Maratos-Flier E, Müller TD, Münzberg H, Pfluger PT, Plum L, Reitman ML, Rahmouni K, Shulman GI, Thomas G, Kahn CR, Ravussin E.** A guide to analysis of mouse energy metabolism. *Nat Met* 2011; 9(1): 57-63
<http://www.ncbi.nlm.nih.gov/pubmed/22205519>
- **Verbaeys I, Tolle V, Swennen Q, Zizzari P, Buyse J, Epelbaum J, Cokelaere.** Scheduled feeding results in adipogenesis and increased acylated ghrelin. *Am Journal Physiol Endocrinol Metab* 2011; 300(6): e1103-11
<http://www.ncbi.nlm.nih.gov/pubmed/21427412>
- **Vernia S, Heredia M, Criado O, Rodriguez de Cordoba S, Garcia-Roves PM, Cansell C, Denis R, Luquet S, Fougelle F, Ferre P, Sanz P.** Laforin, a dual specificity phosphatase involved in lafora disease, regulates insulin response and whole-body energy balance in mice. *Hum Mol Genet* 2011; 20(13): 2571-84
<http://www.ncbi.nlm.nih.gov/pubmed/21493628>
- **Veyrat-Durebex C, Poher AL, Caillon A, Montet X, Rohner-Jeanrenaud F.** Alterations in lipid metabolism and thermogenesis with emergence of brown adipocytes in white adipose tissue in diet-induced obesity-resistant Lou/C rats. *Am J Physiol Endocrinol Metab* 2011; 300: 1146-57
<http://www.ncbi.nlm.nih.gov/pubmed/21406614>
- **Wellner N, Tsuboi K, Madsen AN, Holst B, Diep TA, Nakao M, Tokumura A, Burns MP, Deutsch DG, Ueda N, Hansen HS.** Studies on the anorectic effect of N-acylphosphatidylethanolamine and phosphatidylethanolamine in mice. *Biochim Biophys Acta* 2011; 1811(9): 508-12
<http://www.ncbi.nlm.nih.gov/pubmed/21723414>
- **Wong KE, Kong J, Zhang W, Szeto FL, Ye H, Deb DK, Brady MJ, Chun Li Y.** Targeted expression of human vitamin D receptor in adipocytes decreases energy expenditure and induces obesity in mice. *J Biol Chem* 2011; 286(39): 33804-10
<http://www.ncbi.nlm.nih.gov/pubmed/21840998>
- **Zizzari P, Hassouna R, Longchamps R, Epelbaum J, Tolle V.** Meal anticipatory rise in acylated ghrelin at dark onset is blunted after long-term fasting in rats. *J Neuroendocrinol* 2011; 23: 804-14
<http://www.ncbi.nlm.nih.gov/pubmed/21722214>

2010

- **Baghbanzadeh A, Modirsaneie M, Emam G, Hajinezhad M.** Microhandling of vesicular glutamate uptake modulate feeding in broilers. *J Anim Physiol Anim Nutr* 2010; 94(1): 74-77
<http://www.ncbi.nlm.nih.gov/pubmed/19175460>
- **Boy J, Schmidt T, Schumann U, Grasshoff U, Unser S, Holzmann C, Schmitt I, Karl T, Laccone F, Wolburg H, Ibrahim S, Riess O.** A transgenic mouse model of spinocerebellar ataxia type 3 resembling late disease onset and gender-specific instability of CAG repeats. *Neurobiol Dis* 2010; 37(2): 284-93
<http://www.ncbi.nlm.nih.gov/pubmed/19699305>
- **Czyzyk TA, Nogueiras R, Lockwood JF, McKinzie JH, Coskun T, Pintar JE, Hammond C, Tschöp MH, Statnick MA.** K-Opioid receptors control the metabolic response to a high-energy diet in mice. *FASEB J* 2010; 24: 1151-9
<http://www.ncbi.nlm.nih.gov/pubmed/19917675>
- **Dietrich MO, Antunes C, Geliang G, Liu ZW, Borok E, Nie Y, Xu AW, Souza DO, Gao Q, Diano S, Gao XB, Horvath TL.** Agrp neurons mediate Sirt1's action on the melanocortin system and energy balance: roles for Sirt1 in neuronal firing and synaptic plasticity. *J Neurosci* 2010; 30(35): 11815-25
<http://www.ncbi.nlm.nih.gov/pubmed/20810901>
- **Glier MB, Pissios P, Babich SL, Macdonald MLE, Hayden MR, Maratos-Flier E, Gibson WT.** The metabolic phenotype of SCD1-deficient mice is independent of melanin-concentrating hormone. *Peptides* 2010; 31(1): 123-9
<http://www.ncbi.nlm.nih.gov/pubmed/19883709>
- **Guo K, Yu YH, Hou J, Zhang Y.** Chronic leucine supplementation improves glycemic control in etiologically distinct mouse models of obesity and diabetes mellitus. *Nutr Metab* 2010; 7: 57
<http://www.ncbi.nlm.nih.gov/pubmed/20624298>
- **Kuhlow D, Florian S, von Figura G, Weimer S, Schulz N, Petzke KJ, Zarse K, Pfeiffer AF, Rudolph KL, Ristow M.** Telomerase deficiency impairs glucose metabolism and insulin secretion. *Aging* 2010; 2(10):650-8
<http://www.ncbi.nlm.nih.gov/pubmed/20876939>
- **Labonté ED, Pfluger PT, Cash JG, Kuhel DG, Roja JC, Magness DP, Jandacek RJ, Tschöp MH, Hui DY.** Postprandial lysophospholipid suppresses hepatic fatty acid oxidation: the molecular link between group 1B phospholipase A2 and diet-induced obesity. *FASEB J* 2010; 24(7): 2516-24
<http://www.ncbi.nlm.nih.gov/pubmed/20215528>
- **Li X, Cope MB, Johnson MS, Smith DL Jr, Nagy TR.** Mild calorie restriction induces fat accumulation in female C57BL/6J mice. *Obesity* 2010; 18(3): 456-462
<http://www.ncbi.nlm.nih.gov/pubmed/19798071>
- **Lin HV, Plum L, Ono H, Gutiérrez-Juárez R, Shanabrough M, Borok E, Horvath TL, Rossetti L, Accili D.** Divergent regulation of energy expenditure and hepatic glucose production by insulin receptor on agouti-related protein and POMC neurons. *Diabetes* 2010; 59(2): 337-46
<http://www.ncbi.nlm.nih.gov/pubmed/19933998>

- **Mauer J, Chaurasia B, Plum L, Quast T, Hampel B, Blüher M, Kolanus W, Kahn CR, Brüning JC.** Myeloid cell-restricted insulin receptor deficiency protects against obesity-induced inflammation and systemic insulin resistance. *PloS Genet* 2010; 6(5): e1000938
<http://www.ncbi.nlm.nih.gov/pubmed/20463885>
- **Miesel A, Müller H, Thermann M, Heidbreder M, Dominiak P, Raasch W.** Overfeeding-induced obesity in spontaneously hypertensive rats: an animal model of the human metabolic syndrome. *Ann Nutr Metab* 2010; 56(2): 127-42
<http://www.ncbi.nlm.nih.gov/pubmed/20134158>
- **Ngai YF, Quong WL, Glier MB, Glavas MM, Babich SL, Innis SM, Kieffer TJ, Gibson WT.** Ldlr^{-/-} mice display decreased susceptibility to Western-type diet-induced obesity due to increased thermogenesis. *Endocrinology* 2010; 151(11): 5226-36
<http://www.ncbi.nlm.nih.gov/pubmed/20881250>
- **Peter JC, Lecourt AC, Weckering M, Zipfel G, Niehoff ML, Banks WA, Hofbauer KG.** A pharmacologically active monoclonal antibody against the human melanocortin-4 receptor: effectiveness after peripheral and central administration. *J Pharmacol Exp Ther* 2010; 333(2): 478-90
<http://www.ncbi.nlm.nih.gov/pubmed/20118207>
- **Ring LE, Zeltser LM.** Disruption of hypothalamic leptin signaling in mice leads to early-onset obesity, but physiological adaptations in mature animals stabilize adiposity levels. *J Clin Invest* 2010; 120(8): 2931-41
<http://www.ncbi.nlm.nih.gov/pubmed/20592471>
- **Sabio G, Cavanagh-Kyros J, Barrett T, Jung DY, Ko HJ, Ong H, Morel C, Mora A, Reilly J, Kim JK, Davis RJ.** Role of the hypothalamic-pituitary-thyroid axis in metabolic regulation by JNK1. *Genes Dev* 2010; 24(3): 256-64
<http://www.ncbi.nlm.nih.gov/pubmed/20080940>
- **Sassmann A, Offermanns S, Wettschureck N.** Tamoxifen-inducible Cre-mediated recombination in adipocytes. *Genesis* 2010; 48(10): 618-625
<http://www.ncbi.nlm.nih.gov/pubmed/20715175>
- **Schoiswohl G, Schweiger M, Schreiber R, Gorkiewicz G, Preiss-Landl K, Taschler U, Zierler KA, Radner FPW, Eichmann TO, Kienesberger PC, Eder S, Lass A, Haemmerle G, Alsted TJ, Kiens B, Hoefler G, Zechner R, Zimmermann R.** Adipose triglyceride lipase plays a key role in the supply of the working muscle with fatty acids. *J Lipid Res* 2010; 51(3): 490-9
<http://www.ncbi.nlm.nih.gov/pubmed/19965578>
- **Song Y, Altarejos J, Goodarzi MO, Inoue H, Guo X, Berdeaux R, Kim JH, Goode J, Igata M, Paz JC, Hogan MF, Singh PK, Goebel N, Vera L, Miller N, Cui J, Jones MR, CHARGE Consortium, GIANT Consortium, Chen YDI, Taylor KD, Hsueh WA, Rotter JI, Montminy M.** CRT3 links catecholamine signalling to energy balance. *Nature* 2010; 468(7326): 933-9
<http://www.ncbi.nlm.nih.gov/pubmed/21164481>

- **Soon Kang H, Okamoto K, Kim YS, Takeda Y, Bortner CD, Dang H, Wada T, Xie W, Yang XP, Liao G, Jetten AM.** Nuclear orphan receptor TAK1/TR4-deficient mice are protected against obesity-linked inflammation, hepatic steatosis, and insulin resistance. *Diabetes* 2010; 60(1): 177-88
<http://www.ncbi.nlm.nih.gov/pubmed/20864514>
- **Sutanto MM, Ferguson KK, Sakuma H, Ye H, Brady MJ, Cohen RN.** The silencing mediator of retinoid and thyroid hormone receptors (SMRT) regulates adipose tissue accumulation and adipocyte insulin sensitivity in vivo. *J Biol Chem* 2010; 285(24): 18485-95
<http://www.ncbi.nlm.nih.gov/pubmed/20371609>
- **Urbach YK, Raber KA, von Hörsten S.** Automated physio-behavioral intra-home-cage phenotyping in SCA17 transgenic rats facilitates multidimensional recording of various activity pattern. *Lettre de la Société Francophone de Chronobiologie* 2010; 41(1):29
- **Ye R, Jung DY, Jun JY, Li J, Luo S, Ko HJ, Kim JK, Lee AS.** Grp78 heterozygosity promotes adaptive unfolded protein response and attenuates diet-induced obesity and insulin resistance. *Diabetes* 2010; 59(1): 6-16
<http://www.ncbi.nlm.nih.gov/pubmed/19808896>

2009

- **Caton SJ, Yinglong B, Burget L, Spangler LJ, Tschöp MH, Bidlingmaier M.** Low-carbohydrate high-fat diets: regulation of energy balance and body weight regain in rats. *Obesity* 2009; 17(2): 283-9
<http://www.ncbi.nlm.nih.gov/pubmed/19039316>
- **Danilova OV, Tai AK, Mele DA, Neinborn M, Leiter AB, Greenberg AS, Perfield JW, DeFuria J, Singru PS, Lechan RM, Huber BT.** Neurogenin 3-specific dipeptidyl peptidase-2 deficiency causes impaired glucose tolerance, insulin resistance, and visceral obesity. *Endocrinology* 2009; 150(12): 5240-8
<http://www.ncbi.nlm.nih.gov/pubmed/19819973>
- **Day JW, Ottaway N, Patterson JT, Gelfanov V, Smiley D, Gidda J, Findeisen H, Bruemmer D, Drucker DJ, Chaudhary N, Holland J, Hembree J, Abplanalp W, Grant E, Ruehl J, Wilson H, Kirchner H, Lockie SH, Hofmann S, Woods SC, Nogueiras R, Pfluger PT, Perez-Tilve D, DiMarchi R, Tschöp MH.** A new glucagon and GLP-1 co-agonist eliminates obesity in rodents. *Nat Chem Biol* 2009; 5(10): 749-57
<http://www.ncbi.nlm.nih.gov/pubmed/19597507>
- **Edelsbrunner ME, Herzog H, Holzer P.** Evidence from knockout mice that peptide YY and neuropeptide Y enforce murine locomotion, exploration and ingestive behaviour in a circadian cycle- and gender-dependent manner. *Behav Brain Res* 2009; 203(1): 97-107
<http://www.ncbi.nlm.nih.gov/pubmed/19397935>
- **Fischer J, Koch L, Emmerling C, Vierkotten J, Peters T, Brüning JC, Rüdter U.** Inactivation of the FTO gene protects from obesity. *Nature* 2009; 458(7240): 894-8
<http://www.ncbi.nlm.nih.gov/pubmed/19234441>
- **Kilpeläinen M, Riikonen J, Vlasova MA, Huotari A, Lehto VP, Salonen J, Herzig KH, Järvinen K.** In vivo delivery of a peptide, ghrelin antagonist, with mesoporous silicon microparticles. *J Control Release* 2009; 137(2): 166-70

<http://www.ncbi.nlm.nih.gov/pubmed/19345247>

- **Langmesser S, Franken P, Feil S, Emmenegger Y, Albrecht U, Feil R.** cGMP-dependent protein kinase type 1 is implicated in the regulation of the timing and quality of sleep and wakefulness. *PLoS One* 2009; 4(1): 1-13
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2617781/>
- **Miinalainen IJ, Schmitz W, Huotari A, Autio KJ, Soinen R, Ver Loren van Themaat E, Baes M, Herzig KH, Conzelmann E, Hiltunen K.** Mitochondrial 2,4-dienoyl-CoA reductase deficiency in mice results in severe hypoglycemia with stress intolerance and unimpaired ketogenesis. *PLoS Genet* 2009; 5(7): e1000543
<http://www.ncbi.nlm.nih.gov/pubmed/19578400>
- **Nogueiras R, Perez-Tilve D, Veyrat-Durebex C, Morgan DA, Varela L, Haynes WG, Patterson JT, Disse E, Pfluger PT, Lopez M, Woods SC, DiMarchi R, Dieguez C, Rahmouni K, Rohner-Jeanraud F, Tschöp MH.** Direct control of peripheral lipid deposition by CNS GLP-1 receptor signaling is mediated by the sympathetic nervous system and blunted in diet-induced obesity. *J Neurosci* 2009; 29(18): 5916-25
<http://www.ncbi.nlm.nih.gov/pubmed/19420258>
- **Plum L, Lin HV, Dutia R, Tanaka J, Aizawa KS, Matsumoto M, Kim AJ, Cawley NX, Paik JH, Loh YP, DePinho RA, Wardlaw SL, Accili D.** The obesity susceptibility gene Carboxypeptidase E links FoxO1 signaling in hypothalamic pro-opiomelanocortin neurons with regulation of food intake. *Nat Med* 2009; 15(10): 1195-201
<http://www.ncbi.nlm.nih.gov/pubmed/19767734>
- **Stratigopoulos G, LeDuc CA, Matsuoka N, Gutman R, Rausch R, Robertson SA, Myers MG, Chung WK, Chua SC, Leibel RL.** Functional consequences of the human leptin receptor (LEPR) Q223R transversion. *Obesity* 2009; 17(1): 126-35
<http://www.ncbi.nlm.nih.gov/pubmed/18997673>
- **Stylopoulos N, Hoppin AG, Kaplan LM.** Roux-en-Y gastric bypass enhances energy expenditure and extends lifespan in diet-induced obese rats. *Obesity* 2009; 17(10): 1839-47
<http://www.ncbi.nlm.nih.gov/pubmed/19556976>
- **Tasan RO, Lin S, Hetzenauer A, Singewald N, Herzog H, Sperk G.** Increased novelty-induced motor activity and reduced depression-like behavior in neuropeptide Y (NPY)-Y4 receptor knockout mice. *Neuroscience* 2009; 158(4): 1717-1730
<http://www.ncbi.nlm.nih.gov/pubmed/19121371>
- **Wong KE, Szeto FL, Zhang W, Ye H, Kong J, Zhang Z, Sun XJ, Li YC.** Involvement of the vitamin D receptor in energy metabolism: regulation of uncoupling proteins. *Am J Physiol Endocrinol Metab* 2009; 296(4): e820-8
<http://www.ncbi.nlm.nih.gov/pubmed/19176352>

2008

- **Banks A, Kon N, Knight C, Matsumoto M, Gutiérrez-Juárez R, Rossetti L, Gu W, Accili D.** SirT1 gain of function increases energy efficiency and prevents diabetes in mice. *Cell Metab* 2008; 8(4): 333-41
<http://www.ncbi.nlm.nih.gov/pubmed/18840364>

- **Bode FJ, Stephan M, Suhling H, Pabst R, Straub RH, Raber KA, Bonin M, Nguyen HP, Riess O, Bauer A, Sjoberg C, Petersén A, von Hörsten S.** Sex differences in a transgenic rat model of Huntington's disease: decreased 17 β -estradiol levels correlate with reduced numbers of DARPP32+ neurons in males. *Hum Mol Genet* 2008; 17(17): 2595-609
<http://www.ncbi.nlm.nih.gov/pubmed/18502785>
- **Choi Y, Li C, Hartzell, DL, Little DE, Della-Fera MA, Baile CA.** ICV leptin effects on spontaneous physical activity and feeding behavior in rats. *Behav Brain Res* 2008; 188: 100-8
<http://www.ncbi.nlm.nih.gov/pubmed/18063139>
- **Foryst-Ludwig A, Clemenz M, Hohmann S, Hartge M, Sprang C, Frost N, Krikov M, Bhanot S, Barros R, Morani A, Gustafsson JA, Unger T, Kintscher U.** Metabolic actions of estrogen receptor beta (Erb) are mediated by a negative cross-talk with PPAR γ . *PLoS Genet* 2008; 4(6): e1000108
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2432036/>
- **Hofmann SM, Perez-Tilve D, Greer TM, Coburn BA, Grant E, Basford JE, Tschöp MH, Hui DY.** Defective Lipid Delivery Modulates Glucose Tolerance and Metabolic Response to Diet in Apolipoprotein E-Deficient Mice. *Diabetes* 2008; 57(1): 5-12
<http://www.ncbi.nlm.nih.gov/pubmed/17914034>
- **Jayasooriya AP, Mathai ML, Walker LL, Begg DP, Denton DA, Cameron-Smith D, Egan GF, McKinley MJ, Rodger PD, Sinclair AJ, Wark JD, Weisinger HS, Jois M, Weisinger RS.** Mice lacking angiotensin-converting enzyme have increased energy expenditure, with reduced fat mass and improved glucose clearance. *Proc Nat Acad Sci USA* 2008; 105: 6531-6
<http://www.ncbi.nlm.nih.gov/pubmed/18443281>
- **Painsipp E, Wultsch T, Edelsbrunner ME, Tasan RO, Singewald N, Herzog H, Holzer P.** Reduced anxiety-like and depression-related behavior in neuropeptide Y Y4 receptor knockout mice. *Genes Brain Behav* 2008; 7(5): 532-42
<http://www.ncbi.nlm.nih.gov/pubmed/18221379>
- **Pfluger PT, Herranz D, Velasco S, Serrano M, Tschöp MH.** Sirt1 protects against high-fat diet-induced metabolic damage. *Proc Nat Acad Sci* 2008; 105(28): 9793-8
<http://www.ncbi.nlm.nih.gov/pubmed/18599449>
- **Pfluger PT, Kirchner H, Günnel S, Schrott B, Perez-Tilve D, Fu S, Benoit SC, Horvath T, Joost HG, Wortley KE, Sleeman MW, Tschöp MH.** Simultaneous deletion of ghrelin and its receptor increases motor activity and energy expenditure. *Am J Physiol Gastrointest Liver Physiol* 2008; 294(3): G610-8
<http://www.ncbi.nlm.nih.gov/pubmed/18048479>
- **Pilvi TK, Seppänen-Laakso T, Simolin H, Finckenberg P, Huotari A, Herzig KH, Korpela R, Oresic M, Mervaala EM.** Metabolic changes in fatty liver can be modified by dietary protein and calcium during energy restriction. *World J Gastroenterol* 2008; 14(28): 4462-72
<http://www.ncbi.nlm.nih.gov/pubmed/18680224>
- **Reed JA, Benoit SC, Pfluger PT, Tschöp MH, D'Alessio DA, Seeley RJ.** Mice with chronically increased circulating ghrelin develop age-related glucose intolerance. *Am J Physiol Endocrinol Metab* 2008; 294: e752-60
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3632576/>

- **Schmidt S, Richter M, Montag D, Sartorius T, Gawlik V, Hennige AM, Scherneck S, Himmelbauer H, Lutz SZ, Augustin R, Kluge R, Ruth P, Joost HG, Schürmann A.** Neuronal functions, feeding behavior, and energy balance in Slc2a3^{+/-} mice. *American journal of physiology. Endocrinol Metab* 2008; 295(5): e1084-94
<http://www.ncbi.nlm.nih.gov/pubmed/18780771>
- **Somm E, Schwitzgebel VM, Vauthay DM, Camm EJ, Chen CY, Giacobino JP, Sizonenko SV, Aubert ML, Hüppi PS.** Prenatal nicotine exposure alters early pancreatic islet and adipose tissue development with consequences on the control of body weight and glucose metabolism later in life. *Endocrinology* 2008; 149(12): 6289-99
<http://www.ncbi.nlm.nih.gov/pubmed/18687784>
- **Strassburg S, Anker SD, Castaneda TR, Burget L, Perez-Tilve D, Pfluger PT, Nogueiras R, Halem H, Dong JZ, Culler MD, Datta R, Tschöp MH.** Long-term effects of ghrelin and ghrelin receptor agonists on energy balance in rats. *Am J Physiol Endocrinol Metab* 2008; 295(1): e78-84
<http://www.ncbi.nlm.nih.gov/pubmed/18460598>

2007

- **Cho YR, Kim HJ, Park SY, Ko HJ, Hong EG, Higashimori T, Zhang Z, Jung DY, Ola MS, Lanoue KF, Leiter EH, Kim JK.** Hyperglycemia, maturity-onset obesity, and insulin resistance in NONcNZO10/LtJ males, a new mouse model of type 2 diabetes. *Am J Physiol Endocrinol Metab* 2007; 293(1): e327-36
<http://www.ncbi.nlm.nih.gov/pubmed/17616608>
- **Choi Y, Li C, Hartzell, DL, Little DE, Della-Fera MA, Baile CA.** ICV leptin effects on spontaneous physical activity and feeding behavior in rats. *Behav Brain Res* 2008; 188(1): 100-8
<http://www.ncbi.nlm.nih.gov/pubmed/18063139>
- **Hamrick MW, Della Fera MA, Choi YH, Hartzell D, Pennington C, Baile CA.** Injections of leptin into rat ventromedial hypothalamus increase adipocyte apoptosis in peripheral fat and in bone marrow. *Cell Tissue Res* 2007; 327(1): 133-41
<http://www.ncbi.nlm.nih.gov/pubmed/17024416>
- **Herling AW, Gossel M, Haschke G, Stengelin S, Kuhlmann J, Müller G, Schmoll D, Kramer W.** CB1 receptor antagonist AVE1625 affects primarily metabolic parameters independently of reduced food intake in Wistar rats. *Am J Physiol Endocrinol Metab* 2007; 293(3): e826-32
<http://www.ncbi.nlm.nih.gov/pubmed/17595216>
- **Hofmann SM, Zhou L, Perez-Tilve D, Greer T, Grant E, Wancata L, Thomas A, Pfluger PT, Basford JE, Gilham D, Herz J, Tschöp MH, Hui DY.** Adipocyte LDL receptor-related protein-1 expression modulates postprandial lipid transport and glucose homeostasis in mice. *J Clin Invest* 2007; 117(11): 3271-82
<http://www.ncbi.nlm.nih.gov/pubmed/17948131>
- **Hong EG, Jung DY, Ko HJ, Zhang Z, Ma Z, Jun JY, Kim JH, Summer AD, Vary TC, Gardner TW, Bronson SK, Kim JK.** Nonobese, insulin-deficient Ins2Akita mice develop type 2 diabetes phenotypes including insulin resistance and cardiac remodeling. *Am J Physiol Endocrinol Metab* 2007; 293(6): 1687-96
<http://www.ncbi.nlm.nih.gov/pubmed/17911348>

- **Nogueiras R, Pfluger P, Tovar S, Myrtha A, Mitchell S, Morris A, Perez-Tilve D, Vazquez MJ, Wiedmer P, Castaneda TR, Dimarchi R, Tschop M, Schürmann A, Joost HG, Williams LM, Langhans W, Dieguez C.** Effects of obestatin on energy balance and growth hormone secretion in rodents. *Endocrinology* 2007; 148(1): 21-6
<http://www.ncbi.nlm.nih.gov/pubmed/17008393>
- **Nogueiras R, Wiedmer P, Perez-Tilve D, Veyrat-Durebex C, Keogh JM, Sutton GM, Pfluger PT, Castaneda TR, Neschen S, Hofmann SM, Howles PN, Morgan DA, Benoit SC, Szanto I, Schrott B, Schürmann A, Joost HG, Hammond C, Hui DY, Woods SC, Rahmouni K, Butler AA, Farooqi IS, O'Rahilly S, Rohner-Jeanrenaud F, Tschöp MH.** The central melanocortin system directly controls peripheral lipid metabolism. *J Clin Invest* 2007; 117(11): 3475-88
<http://www.ncbi.nlm.nih.gov/pubmed/17885689>
- **Nomiyama T, Perez-Tilve D, Ogawa D, Gizard F, Zhao Y, Heywood EB, Jones KL, Kawamori R, Cassis LA, Tschöp MH, Bruemmer D.** Osteopontin mediates obesity-induced adipose tissue macrophage infiltration and insulin resistance in mice. *J Clin Invest* 2007; 117(10): 2877-88
<http://www.ncbi.nlm.nih.gov/pubmed/17823662>
- **Painsipp E, Wultsch T, Shahbazian A, Edelsbrunner M, Kreissl MC, Schirbel A, Bock E, Pabst MA, Thoeringer CK, Huber HP, Holzer P.** Experimental gastritis in mice enhances anxiety in a gender-related manner. *Neuroscience* 2007; 150(3): 522-36
<http://www.ncbi.nlm.nih.gov/pubmed/17945426>
- **Peter JC, Nicholson JR, Heydet D, Lecourt AC, Hoebeke J, Hofbauer KG.** Antibodies against the melanocortin-4 receptor act as inverse agonists in vitro and in vivo. *Am J Physiol Regul Integr Comp Physiol* 2007; 292(6): R2151-8
<http://www.ncbi.nlm.nih.gov/pubmed/17322114>
- **Sakkou M, Wiedmer P, Anlag K, Hamm A, Seuntjens E, Ettwiller L, Tschop MH, Treier M.** A role for brain-specific homeobox factor *bsx* in the control of hyperphagia and locomotory behavior. *Cell Metab* 2007; 5(6): 450-63
<http://www.ncbi.nlm.nih.gov/pubmed/17550780>
- **Stepanyan Z, Kocharyan A, Behrens M, Koebnick C, Pyrski M, Meyerhof W.** Somatostatin, a negative-regulator of central leptin action in the rat hypothalamus. *J Neurochem* 2007; 100(2): 468-78
<http://www.ncbi.nlm.nih.gov/pubmed/17083445>

2006

- **Biebermann H, Castañeda TR, van Landeghem F, von Deimling A, Escher F, Brabant G, Hebebrand J, Hinney A, Tschöp MH, Grüters A, Krude H.** A role for beta-melanocyte-stimulating hormone in human body-weight regulation. *Cell Metab* 2006; 3(2): 141-6
<http://www.ncbi.nlm.nih.gov/pubmed/16459315>
- **Jurgens HS, Schürmann A, Kluge R, Ortman S, Klaus S, Joost HG, Tschöp MH.** Hyperphagia, lower body temperature, and reduced running wheel activity precede development of morbid obesity in New Zealand obese mice. *Physiol Genomics* 2006; 25(2): 234-41
<http://www.ncbi.nlm.nih.gov/pubmed/16614459>

- **Kampe J, Wiedmer P, Pfluger PT, Castaneda TR, Burget L, Mondala H, Kerr J, Liaw C, Oldfield BJ, Tschöp MH, Bagnol D.** Effect of central administration of QRFP(26) peptide on energy balance and characterization of a second QRFP receptor in rat. *Brain Res* 2006; 1119(1): 133-49
<http://www.ncbi.nlm.nih.gov/pubmed/16996040>
- **Sanna B, Brandt EB, Kaiser RA, Pfluger P, Witt SA, Kimball TR, van Rooij E, De Windt LJ, Rothenberg ME, Tschop MH, Benoit SC, Molkentin JD.** Modulatory calcineurin-interacting proteins 1 and 2 function as calcineurin facilitators in vivo. *Proc Natl Acad Sci USA* 2006; 103(19): 7327-32
<http://www.ncbi.nlm.nih.gov/pubmed/16648267>
- **Theander-Carrillo C, Wiedmer P, Cettour-Rose P, Nogueiras R, Perez-Tilve D, Pfluger P, Castaneda TR, Muzzin P, Schurmann A, Szanto I, Tschop MH, Rohner-Jeanrenaud F.** Ghrelin action in the brain controls adipocyte metabolism. *J Clin Invest* 2006; 116(7): 1983-93
<http://www.ncbi.nlm.nih.gov/pubmed/16767221>

2005

- **Della-Fera MA, Choi YH, Hartzell DL, Duan J, Hamrick M, Baile CA.** Sensitivity of ob/ob mice to leptin-induced adipose tissue apoptosis. *Obes Res* 2005; 13(9): 1540-7
<http://www.ncbi.nlm.nih.gov/pubmed/16222056>
- **Fegley D, Gaetani S, Duranti A, Tontini A, Mor M, Tarzia G, Piomelli D.** Characterization of the fatty acid amide hydrolase inhibitor cyclohexyl carbamic acid 3'-carbamoyl-biphenyl-3-yl ester (URB597): effects on anandamide and oleoylethanolamide deactivation. *J Pharmacol Exp Ther* 2005; 313(1): 352-8
<http://www.ncbi.nlm.nih.gov/pubmed/15579492>
- **Fu J, Oveisi F, Gaetani S, Lin E, Piomelli D.** Oleoylethanolamide, an endogenous PPAR-alpha agonist, lowers body weight and hyperlipidemia in obese rats. *Neuropharmacology* 2005; 48(8): 1147-53
<http://www.ncbi.nlm.nih.gov/pubmed/15910890>
- **Morens C, Norregaard P, Receveur JM, van Dijk G, Scheurink AJW.** Effects of MCH and a MCH1-receptor antagonist on (palatable) food and water intake. *Brain Res* 2005; 1062(1-2): 32-8
<http://www.ncbi.nlm.nih.gov/pubmed/16257393>
- **Siegmund S, Vengeliene V, Singer MV, Spanagel R.** Influence of age at drinking onset on long-term ethanol self-administration with deprivation and stress phases. *Alcohol Clin Exp Res* 2005; 29(7): 1139-45
<http://www.ncbi.nlm.nih.gov/pubmed/16046868>
- **Zizzari P, Halem H, Taylor J, Dong JZ, Datta R, Culler MD, Epelbaum J, Bluet-Pajot MT.** Endogenous ghrelin regulates episodic growth hormone (GH) secretion by amplifying GH pulse amplitude: evidence from antagonism of the GH secretagogue-R1a receptor. *Endocrinology* 2005; 146(9): 3836-42
<http://www.ncbi.nlm.nih.gov/pubmed/15919752>

2004

- **Bickel M, Gossel M, Geisen K, Jaehne G, Lang HJ, Rosenburg R, Sandow J.** Analysis of the anorectic efficacy of HMR1426 in rodents and its effects on gastric emptying in rats. *Int J Obes Relat Metab Disord* 2004; 28(2): 211-21
<http://www.ncbi.nlm.nih.gov/pubmed/14663491>
- **Della-Zuana O, Revereault L, Beck-Sickinger A, Monge A, Caignard DH, Fauchère JL, Henlin JM, Audinot V, Boutin JA, Chamorro S, Félétou M, Levens N.** A potent and selective NPY Y5 antagonist reduces food intake but not through blockade of the NPY Y5 receptor. *Int J Obes Relat Metab Disord* 2004; 28(4): 628-39
<http://www.ncbi.nlm.nih.gov/pubmed/14758341>
- **Oveisi F, Gaetani S, Eng KT, Piomelli D.** Oleoylethanolamide inhibits food intake in free-feeding rats after oral administration. *Pharmacol Res* 2004; 49(5): 461-6
<http://www.ncbi.nlm.nih.gov/pubmed/14998556>

2003

- **Fu J, Gaetani S, Oveisi F, Lo Verme J, Serrano A, Rodriguez de Fonseca F, Rosengarth A, Luecke H, Di Giacomo B, Tarzia G, Piomelli D.** Oleoylethanolamide regulates feeding and body weight through activation of the nuclear receptor PPAR-alpha. *Nature* 2003; 425(6953): 90-3
<http://www.ncbi.nlm.nih.gov/pubmed/12955147>
- **Gaetani S, Oveisi F, Piomelli D.** Modulation of meal pattern in the rat by the anorexic lipid mediator oleoylethanolamide. *Neuropsychopharmacology* 2003; 28(7): 1311-16.
<http://www.ncbi.nlm.nih.gov/pubmed/12700681>
- **Kathuria S, Gaetani S, Fegley D, Valiño F, Duranti A, Tontini A, Mor M, Tarzia G, La Rana G, Calignano A, Giustino A, Tattoli M, Palmery M, Cuomo V, Piomelli D.** Modulation of anxiety through blockade of anandamide hydrolysis. *Nat Med* 2003; 9(1): 76-81
<http://www.ncbi.nlm.nih.gov/pubmed/12461523>
- **Vengeliene V, Siegmund S, Singer MV, Sinclair JD, Li TK, Spanagel R.** A comparative study on alcohol-preferring rat lines: effects of deprivation and stress phases on voluntary alcohol intake. *Alcohol Clin Exp Res* 2003; 27(7): 1048-54
<http://www.ncbi.nlm.nih.gov/pubmed/12878910>

2002

- **Bauhofer A, Witte K, Lemmer B, Middeke M, Lorenz W, Celik I.** Quality of life in animals as a new outcome for surgical research: G-CSF as a quality of life improving factor. *Eur Surg Res* 2002; 34(1-2): 22-29
<http://www.ncbi.nlm.nih.gov/pubmed/11867897>
- **Brakebusch C, Seidenbecher C, Asztely F, Rauch U, Matthies H, Meyer H, Krug M, Bockers T, Zhou X, Kreutz M, Montag D, Gundelfinger E, Fassler R.** Brevican-deficient mice display impaired hippocampal CA1 long-term potentiation but show no obvious deficits in learning and memory. *Mol Cell Biol* 2002;22(21): 7417-27
<http://www.ncbi.nlm.nih.gov/pubmed/12370289>



- **Sammut S, Bethus I, Goodall G, Muscat R.** Antidepressant reversal of interferon-alpha-induced anhedonia. *Physiol Behav* 2002; 75: 765-72
<http://www.ncbi.nlm.nih.gov/pubmed/12020742>

2001

- **Sammut S, Goodall G, Muscat R.** Acute interferon-alpha administration modulates sucrose consumption in the rat. *Psychoneuroendocrinology* 2001; 26(3): 261-72
<http://www.ncbi.nlm.nih.gov/pubmed/11166489>

1999

- **Wallerath T, Witte K, Schaefer SC, Schwarz PM, Prellwitz W, Wohlfahrt P, Kleinert H, Lehr HA, Lemmer B, Foerstermann U.** Down-regulation of the expression of endothelial NO synthase is likely to contribute to glucocorticoid-mediated hypertension. *Proc Natl Acad Sci* 1999; 96(23): 13357-62
<http://www.ncbi.nlm.nih.gov/pubmed/10557325>



Service & Warranty

TSE Systems offers a Two (2) Years ALL-IN Premium Warranty with all new products, including:

- 24/7 technical hotline
- Remote maintenance and update function
- On-site visits upon necessity
- Free replacement parts during warranty

After the expiry of the warranty period, TSE Systems offers comprehensive extensions of the warranty or economical maintenance and repair contracts to ensure the continued smooth running of your instruments. Please contact us for further details.